



EDITORIAL

The Geographical and Economical Impact of Scrub Typhus, the Fastest-growing Vector-borne Disease in Korea

Scrub typhus, or Tsutsugamushi disease is a vector transmitted infectious and febrile illness caused by "Orientia tsutsugamushi" bacteria. It is transmitted to humans through larvae bites of different species of trombiculid mites. The habitat of mites is located in low trees and bushes. However this vector can live in many different areas even in sandy and mountain desert [1].

In 1985, infected people appeared again [2]. Since then Scrub typhus incidences have been reported every year. Now it is considered as one of the most prevalent diseases affecting humans Korea especially in the southwestern provinces of the country [3]. Usually, the larvae of mites feed on wild rats and the human gets infected accidentally in a zone of infected mites. The chance of outbreak of this disease depends on the number of contacts between the human and the mites. Hence the habitats of mites and the human activities are the key factors in the *Scrub typhus* prevalence. Human activities include war, farming, camping, urbanization and so on. A study on the epidemiological characteristic of this disease in Korea supports this general notion [4].

Chigger population densities were great in areas with high relative humidity, low temperature, low incident sunlight and a dense substrate vegetative canopy [5]. The relationship between human incidence of scrub typhus and climate should largely reflect the responses of chiggers to the environment [6].

The occurrence of scrub typhus is also related to land uses such as urbanization and developing cornfield and bio-fuel productions and oil palms [7]. The replacement of natural forests with plantations at forest fringe facilitated the establishment of the disease, providing the best habitats for the mite [8]. Malaysia has seen an

increasing incidence of scrub typhus which is directly related with large-scale cultivation of oil palm and rubber [9]. Urbanization means that the habitats of mites and the life style of human are changing. The contact rate of human with wild life is getting higher. We need information of the habitats of mites and the human activities. Also the incidences of the scrub typhus could be also influenced by climate change in Korea [10].

The geographical information is quite important to trace and predict the occurrence of *Scrub typhus* in spatial respect. It is valuable to use the geographic information system (GIS) method to help the analysis of the occurrence trend. It is widely used in the management of vector-borne disease and human health [11–13]. GIS method can be useful to evaluate the habitat of wildlife [14,15]. Recently it is applied to spatial analysis of scrub typhus [16].

The infected persons of *Scrub typhus* increased more than three times during the period 2001–2012. The prevalence of this disease in Korea has been on a steady rise over the past years. As pointed out by WHO, the reason for such high prevalence of *Scrub typhus* is due to climate change. The major methods used to estimate the loss caused by diseases is contingent valuation [CV] method that encompasses all the implicit values. Diener, O'Brien & Gafni [1998] compared research results based on CV while Yen, et. al. [2007] estimated the value of vaccine that blocks the infection of SARS in Taiwan [17,18]. Krupnick, et. al. [2002] analyzed the loss of value caused by death-causing diseases in Ontario, Canada [19]. Alberini, Hunt & Markandya [2006] estimated WTP for value of a statistical life in EU countries such as the UK, Italy and France [20]. The

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

CV method was used for the social loss value of avian influenza, zoonoses in Korea [21–23].

In this issue, Jin et al. have analyzed the spatial pattern of the occurrence of scrub typhus in Korea using the data from the Korea Centers for Disease Control and Prevention. And the correlation with occurrence of scrub typhus and land use change is studied. The authors concluded that the Gangwon Province and Gyeongsangbuk Province show low incidence number all through the year. Some districts have almost identical environmental condition of *Scrub typhus* incidence. The land use change of districts does not affect directly the incidence rate. GIS analysis shows spatial characteristics of *Scrub typhus* [24].

Rhee has adopted the climate change in the incidence of *Scrub typhus* in Korea. This research can be used to construct spatial-temporal model to understand the epidemic *Scrub typhus*. Double Bounded Dichotomous Choice of Contingent Valuation method is used to estimate Willingness to Payment to avoid infection of this disease, through the survey in the patient group and the control group. The younger the age of the family is, the higher the level of awareness on risks caused by climate change, men, the higher the income is, the lower the suggested bid is, the higher WTP is to avoid infection of disease. The means of the amount of WTP are estimated to be 3,689 KRW per month. As people have become increasingly aware of climate change diseases, WTP to avoid infection of *Scrub typhus* has increased accordingly. And the implicit loss of value due to climate change diseases is increasingly becoming higher. Therefore, there should be stronger and more aggressive promotional activities recommended [25].

References

1. World Health Organization. Frequently Asked Questions: Scrub Typhus 2009. Available from: http://www.searo.who.int/LinkFiles/CDS_faq_Scrub_Typhus.pdf.
2. Bang HA, Lee MJ, Lee WC. Comparative research on epidemiological aspects of tsutsugamushi disease (scrub typhus) between Korea and Japan. Japanese journal of infectious diseases 2008 Mar;61(2):148–50. PubMed PMID: 18362409.
3. Korea Centers for Disease Control and Prevention. Current trends of scrub typhus. Communicable Disease Monthly report 2004;15:245–52.
4. Kim S, Kim JS, Lee H. Epidemiological characteristic of Scrub typhus in Korea 2007–2009. Osong Public Health Res Perspect 2010 Dec;1(1):55–60.
5. Clopton RE, Gold RE. Distribution and seasonal and diurnal activity patterns of *Eutrombicula alfreddugesi* (Acari: Trombiculidae) in a forest edge ecosystem. Journal of medical entomology 1993 Jan;30(1):47–53. PubMed PMID: 8433345.
6. Kawamura A, Tanaka H, Takamura A. Tsutsugamushi Disease: an Overview. Tokyo: University of Tokyo Press; 1995.
7. Rockstrom J, Steffen W, Noone K, et al. A safe operating space for humanity. Nature 2009;461(7263):472–5. <http://dx.doi.org/10.1038/461472a>. PubMed PMID: 19779433.
8. Kim BN, Gordillo LF, Kim Y. A model for the transmission dynamics of *Orientia tsutsugamushi* among its natural reservoirs. Journal of theoretical biology 2010 Sep;266(1): 154–61. <http://dx.doi.org/10.1016/j.jtbi.2010.06.015>. PubMed PMID: 20600142.
9. Hoe KB. Environmental change, development and vector-borne disease: Malaysia's experience with filariasis, scrub typhus and dengue. Environment, Development and Sustainability 2008;10(2):209–17. Epub 217.
10. Kim SH, Jang JY. Correlations between climate change-related infectious diseases and meteorological factors in Korea. Journal of Preventive Medicine and Public Health 2010 Sep;43(5):436–44.
11. Estrada-Pena A, Venzal JM. A GIS framework for the assessment of tick impact on human health in a changing climate. Geo-spatial health 2007 May;1(2):157–68. PubMed PMID: 18686241.
12. Lozano-Fuentes S, Elizondo-Quiroga D, Farfan-Ale JA, et al. Use of Google Earth to strengthen public health capacity and facilitate management of vector-borne diseases in resource-poor environments. Bulletin of the World Health Organization 2008 Sep;86(9):718–25. PubMed PMID: 18797648; PubMed Central PMCID: PMC2649496.
13. Rogers DJ, Randolph SE, Snow RW, et al. Satellite imagery in the study and forecast of malaria. Nature 2002 Feb; 415(6872):710–5. <http://dx.doi.org/10.1038/415710a>. PubMed PMID: 11832960; PubMed Central PMCID: PMC3160466.
14. Gerrard R, Stine P, Church R, et al. Habitat evaluation using GIS – a case study applied to the San Joaquin Kit Fox. Landscape Urban Plan 2001;52(4):239–55. [http://dx.doi.org/10.1016/S0169-2046\(00\)00119-5](http://dx.doi.org/10.1016/S0169-2046(00)00119-5). PubMed PMID: WOS: 000166829500003.
15. Jose MCP, Itami RM. GIS-based habitat modeling using logistic multiple regression: a study of the mt. graham red squirrel. Photogrammetric Engineering and Remote Sensing 1991;57(11). 147501486. Epub 1486.
16. Kuo CC, Huang JL, Ko CY, Lee PF, Wang HC. Spatial analysis of scrub typhus infection and its association with environmental and socioeconomic factors in Taiwan. Acta tropica 2011 Oct–Nov;120(1–2):52–8. <http://dx.doi.org/10.1016/j.actatropica.2011.05.018>. PubMed PMID: 21703220.
17. Diener A, O'Brien B, Gafni A. Health care contingent valuation studies: a review and classification of the literature. Health Economics 1998;7(4):313–26.
18. Yen ZS, Chang CJ, Chen SY, et al. How much would you be willing to pay for preventing a new dangerous infectious disease: a willingness-to-pay 9 study in medical personnel working in emergency department. American Journal of Infection Control 2007;8:213–40.
19. Krupnick A, Alberini A, Cropper M, et al. Age, health and the willingness to pay for mortality risk reductions: a contingent valuation survey of Ontario residents. Journal of Risk and Uncertainty 2002;24(2):161–86.
20. Alberini A, Hunt A, Markandya A. Willingness to pay to reduce mortality risks: evidence from a three-country contingent valuation study. Environmental and Resource Economics 2006;33(2):251–64.
21. Rhee HC, Lim HS. The economic cost of ZOONOSSES: in the case of avian influenza. Korea J Health Economics and Policy; 2007:19–40 [in Korean].
22. Kim TJ, Rhee HC, Lim H.S. Socio-economic Assessment of Loss Brucella, KNIH; 2007 [in Korean].
23. Rhee HC, Kim JH, Lee E. Estimation of willingness to pay for avoiding infection of tsutsugamushi disease in Korea: using contingent valuation method. The Korean Journal of Health Economics and Policy 2009;15(1):131–55 [in Korean].
24. Jin HS, Chu C, Han DY. Spatial Distribution Analysis of Scrub typhus in Korea. Osong Public Health Res Perspect 2013 Feb; 4(1):4–15.

25. Rhee H-C. Willingness to Payment for avoiding infection of Climate Change Diseases: In case of Tsutsugamushi Disease. *Osong Public Health Res Perspect* 2013 Feb;4(1): 16–20.

Hae-Wol Cho, Editor-in-Chief, Dean, Professor*
Osong Public Health and Research Perspectives,
Osong, Korea

College of Health Industry, Eulji University,
Seongnam, Korea

College of Medicine, Eulji University,
Daejeon, Korea

*Corresponding author.

E-mail: hwcho@eulji.ac.kr

Chaeshin Chu, PhD, Managing Editor*
Osong Public Health and Research Perspectives,
Osong, Korea

*Corresponding author.

E-mail: cchu@cdc.go.kr